

NON-PRODUCTIVE DUALITY PROPERTIES OF TOPOLOGICAL GROUPS

MASASI HIGASIKAWA

ABSTRACT. We address two properties for Abelian topological groups: “every closed subgroup is dually closed” and “every closed subgroup is dually embedded.” We exhibit a pair of topological groups such that each has both of the properties and the product has neither, which refutes a remark of N. Noble.

These examples are the additive group of integers topologized with respect to a convergent sequence as investigated by E.G. Zelenyuk and I.V. Protasov. The proof for the product relies on a theorem on exponential Diophantine equations.

1. INTRODUCTION

Several duality properties are well-established for locally-compact Abelian groups. Although some are shared beyond the class ([3], [4], [5], [11]), the boundaries may be rather vague. Our concern is for two of them: “every closed subgroup is dually closed” and “every closed subgroup is dually embedded.” We denote the former by $X(1)$ and the latter by $X(2)$ after [3].

Some definitions and conventions follow. All topological groups treated here are Hausdorff and Abelian, and a character is a continuous homomorphism into the torus $\mathbf{T} = \mathbf{R}/\mathbf{Z}$, unless otherwise stated. A subgroup H of a topological group G is dually closed if for each $g \in G \setminus H$, there exists a character χ of G that separates H and g , i.e., χ is identically zero on H and $\chi(g) \neq 0$. We say that H is dually embedded if each character of H extends to one of G .

That a locally-compact group has $X(1)$ and $X(2)$ is part of the celebrated Pontryagin-van Kampen duality theorem. S. Kaplan [4] shows the same for a product of locally-compact groups. Erroneously referring to the proofs, N. Noble claims in Section 3 of [5] that $X(1)$ and $X(2)$ are each preserved under arbitrary products.

Remark 1.1. Kaplan’s arguments only show that a product $\prod_{i \in J} G_i$ of topological groups has $X(1)$ ($X(2)$, respectively) whenever each subproduct $\prod_{i \in J} G_i$ does for finite $J \subseteq I$. So the problem is reduced to the preservation under finite products, which is out of question for locally-compact groups.

We exhibit a counterexample to Noble’s assertion above. Let $\mathbf{Z}\{2^n\}$ denote the additive group of integers with the strongest group topology such that 2^n converges to 0, and $\mathbf{Z}\{3^n\}$ similarly (E.G. Zelenyuk and I.V. Protasov [12]). Each has both

Date: October 26, 2000.

2000 Mathematics Subject Classification. Primary 22A05, 43A40; Secondary 11D61, 11Z05.

Key words and phrases. Topological group, character, duality, sequential topology, exponential Diophantine equation.

Work partly supported by JSPS Research Fellowships for Young Scientists and a Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture of Japan.

of $X(1)$ and $X(2)$ (Proposition 2.2). We show that the product $\mathbf{Z}\{2^n\} \times \mathbf{Z}\{3^n\}$ has neither (Theorem 2.4) as to the diagonal. The proof is dependent on the fact that the diagonal is closed and discrete (Theorem 4.4), which we get by invoking a theorem on exponential Diophantine equations (cf. [8]).

In Section 2, we investigate some basic properties of the topological group $\mathbf{Z}\{p^n\}$. Section 3 is for number theory necessary for the rest of the argument. The proof of the non-preservation of $X(1)$ and $X(2)$ is completed in Section 4. In Section 5, we construct another counterexample with metrics which have close relation to certain family of exponential Diophantine equations.

2. GROUP TOPOLOGIES WITH A SEQUENCE CONVERGENT

Let G be a group, Abelian or not, and $\langle a_n : n \in \mathbf{N} \rangle$ a sequence in G . Then there exists the strongest group topology on G such that $\langle a_n : n \in \mathbf{N} \rangle$ converges to the neutral element. We denote by $G\{a_n\}$ the topological group G with this topology, which need not be Hausdorff. Zelenyuk and Protasov [12] investigates such topological groups determined by a convergent filter in Abelian and Hausdorff case; unfortunately their recently published monograph [7] have not been available for the author.

The strongest topology thus defined has a useful characterization as follows.

Proposition 2.1. *Suppose that G and $\langle a_n : n \in \mathbf{N} \rangle$ are as above. Let H be a topological group and $f : G \rightarrow H$ a homomorphism between abstract groups. Then f is continuous if and only if $f(a_n) \rightarrow 1$ in H .* $\square$

Our main concern is for topological groups of the form $\mathbf{Z}\{p^n\}$ with p prime. Their closed subgroups and characters are explicitly described in [12].

Proposition 2.2. (1) *The closed subgroups of $\mathbf{Z}\{p^n\}$ are $p^m\mathbf{Z}$ ($m = 0, 1, \dots$) and $\{0\}$.*
 (2) *The character group of $\mathbf{Z}\{p^n\}$ is (identified with) $\mathbf{Z}[1/p]/\mathbf{R} \subset \mathbf{T}$, where $\mathbf{Z}[1/p]$ is the ring generated by $1/p$ over $\mathbf{Z}$.*
 (3) *Both of $X(1)$ and $X(2)$ hold for $\mathbf{Z}\{p^n\}$.* $\square$

Let p and q be distinct primes. The characters of the product $\mathbf{Z}\{p^n\} \times \mathbf{Z}\{q^n\}$ are induced by those of the factors, and hence we see how they behave to the diagonal $\Delta = \{\langle u, u \rangle : u \in \mathbf{Z}\}$.

Lemma 2.3. (1) *The diagonal and each element lying outside cannot be separated by the characters.*
 (2) *The group of characters of Δ extendable to the whole product is a proper subgroup of $\mathbf{T}$.*

Proof. Each character $\chi \in \mathbf{Z}\{p^n\} \times \mathbf{Z}\{q^n\}$ is of form $\chi(u, v) = \chi(1, 0)u + \chi(0, 1)v$ with $\chi(1, 0) \in \mathbf{Z}[1/p]/\mathbf{R}$ and $\chi(0, 1) \in \mathbf{Z}[1/q]/\mathbf{R}$. $\square$

Therefore assuming that the diagonal is closed and discrete (Theorem 4.4), we have the desired non-preservation result.

Theorem 2.4. *The product $\mathbf{Z}\{p^n\} \times \mathbf{Z}\{q^n\}$ has neither $X(1)$ nor $X(2)$.* $\square$

Remark 2.5. By [12, Theorem 6], $\mathbf{Z}\{p^n\}$ is not Fréchet-Urysohn, so not metrizable a fortiori. We observe that it is not α_4 (cf. [1], [2]) due to [12, Lemma 3]; Nyikos [6] proves that Fréchet-Urysohn group is α_4 .

3. S -UNIT EQUATIONS

We recall a finiteness theorem for exponential Diophantine equations, a special case of [8, Ch. V, Theorem 2A]. Let S be a finite set of primes. A rational number is said to be an S -unit if it belongs to the multiplicative group generated by $S \cup \{-1\}$. The set of S -units is denoted by U_S .

Theorem 3.1. *Up to scalar multiplications, the equation $x_1 + \cdots + x_k = 0$ has only finitely many solutions $\langle x_1, \dots, x_k \rangle$ in S -units whose non-trivial subsums do not vanish.* $\square$

As a corollary, we have finiteness for subsums as to a kind of S -unit equation, which we need in the next section.

Lemma 3.2. *Suppose that S and T are disjoint finite sets of primes. Let the tuple $\langle x_1, \dots, x_k, y_1, \dots, y_l \rangle$ runs through the solutions of the equation $x_1 + \cdots + x_k = y_1 + \cdots + y_l$ with $x_i \in U_S \cup \{0\}$ and $y_j \in U_T \cup \{0\}$ for $1 \leq i \leq k, 1 \leq j \leq l$. Then the sum $x_1 + \cdots + x_k$ has only finitely many values.*

Proof. For notational convenience, we consider the equation $x_1 + \cdots + x_k + x_{k+1} + \cdots + x_{k+l} = 0$ replacing each y_j with $-x_{k+j}$. For $I \subseteq \{i : 1 \leq i \leq k+l\}$, we say a tuple $\langle x_i : i \in I \rangle$ to be an I -solution and $\sum_{1 \leq i \leq k, i \in I} x_i$ an I -admissible sum if $\sum_{i \in I} x_i = 0$ with $\{x_i : 1 \leq i \leq k, i \in I\} \subseteq U_S \cup \{0\}$ and $\{x_i : k+1 \leq i \leq k+l, i \in I\} \subseteq U_T \cup \{0\}$. We shall show that there are only finitely many I -admissible sums.

Note that if I is contained in $\{i : 1 \leq i \leq k\}$ or $\{i : k+1 \leq i \leq k+l\}$, which we call a degenerate case, then the only I -admissible sum is 0.

We show the finiteness by the induction on the size of I . If $|I| = 1$, then it is degenerate.

Assume $|I| \geq 2$ and it is not degenerate. Then the equivalence among I -solutions up to scalar multiplications is just the identity. An I -admissible sum is either as to an I -solution whose non-trivial subsums do not vanish or of the form $u + u'$ such that u is a J -admissible sum and u' is an $I \setminus J$ -admissible sum for a non-trivial subset J of I . By the theorem and by the induction hypothesis respectively, there are only finitely many such sums. $\square$

4. REDUCTION TO NUMBER THEORY

We investigate the topological properties of $\mathbf{Z}\{p^n\} \times \mathbf{Z}\{q^n\}$ so that the result in the previous section may be applicable.

Lemma 4.1. *For topological groups of the form $G\{a_n\}, H\{b_n\}$, which need not be Abelian or Hausdorff, the product topology $G\{a_n\} \times H\{b_n\}$ is the same as $(G \times H)\{c_n\}$, where $c_{2n} = \langle a_n, 1 \rangle, c_{2n+1} = \langle 1, b_n \rangle$.*

Proof. It is straightforward that $c_n \rightarrow \langle 1, 1 \rangle$ in $G\{a_n\} \times H\{b_n\}$. So the product has at most as strong topology as that determined by $\langle c_n : n \in \mathbf{N} \rangle$. Hence it is sufficient to show that the identity map $\text{id} : G\{a_n\} \times H\{b_n\} \rightarrow (G \times H)\{c_n\}$ is continuous.

In general, the product topology of topological groups G and H has following characterization. Let $i : G \rightarrow G \times H$ and $j : H \rightarrow G \times H$ be natural injections: $i(g) = \langle g, 1 \rangle, j(h) = \langle 1, h \rangle$. Suppose that K is a topological group and $f : G \times H \rightarrow K$ is a homomorphism between abstract groups. Then f is continuous if and only if so are both of $f \circ i$ and $f \circ j$.

Due to Proposition 2.1, $\text{id} \circ i : G\{a_n\} \rightarrow (G \times H)\{c_n\}$ and $\text{id} \circ j : H\{b_n\} \rightarrow (G \times H)\{c_n\}$ are continuous. So we are done. $\square$

We depend on the following two results in [12] concerning Hausdorff abelian topological groups of the form $G\{a_n\}$.

Lemma 4.2 ([12, Lemma 2]). *If $g_m \rightarrow 0$ in $G\{a_n\}$, then there exists a positive integer k such that $g_m \in \{x_1 + \cdots + x_k : (\forall i)(x_i \in \{\pm a_n : n \in \mathbf{N}\} \cup \{0\})\}$ for sufficiently large m .* $\square$

Theorem 4.3 ([12, Theorem 7]). *The topology of $G\{a_n\}$ is sequential.* $\square$

Now the last piece of the proof follows. Note that every discrete subgroup of a Hausdorff topological group is closed.

Theorem 4.4. *The diagonal Δ is discrete (and closed) in $\mathbf{Z}\{p^n\} \times \mathbf{Z}\{q^n\}$.*

Proof. Applying Theorem 4.3 to $\mathbf{Z}\{p^n\} \times \mathbf{Z}\{q^n\}$ (Lemma 4.1), we observe that the product is also sequential. So it is sufficient to show that Δ has no nontrivial convergent sequence.

Suppose that $\langle \langle u_i, u_i \rangle : i \in \mathbf{N} \rangle$ is a sequence in Δ converging to $\langle 0, 0 \rangle$. By Lemma 4.2, there exists k such that u_i is a sum of less than k numbers in $\{\pm 2^n : n \in \mathbf{N}\}$ and in $\{\pm 3^n : n \in \mathbf{N}\}$, respectively, for sufficiently large i . Due to Lemma 3.2, there are only finitely many such sums. Therefore u_i is eventually equal to 0. $\square$

Remark 4.5. Sequentiality need not be preserved under direct products even for topological groups ([10], [9]).

5. METRIC COUNTEREXAMPLE

Proposition 2.2 holds not only for $\mathbf{Z}\{p^n\}$ but also for a group topology such that each $p^n \mathbf{Z}$ is closed and $p^n \rightarrow 0$. Among those topologies the strongest is that of $\mathbf{Z}\{p^n\}$ and the weakest is the p -adic topology. They have the same closed (open) subgroups and characters.

If $\mathcal{T}$ is such a topology and $\mathcal{U}$ is one for another prime $q \neq p$, then Lemma 2.3 is true for $\langle \mathbf{Z}, \mathcal{T} \rangle \times \langle \mathbf{Z}, \mathcal{U} \rangle$ as well. So we have another counterexample, which we shall construct in this section, provided that the diagonal is discrete.

Suppose that $\delta : \{p^n : n \in \mathbf{N}\} \rightarrow \mathbf{R}_{>0}$ is a non-increasing function with $\delta(p^n) \rightarrow 0$. We define a function $\|\cdot\|_\delta : \mathbf{Z} \rightarrow \mathbf{R}$ by

$$\|u\|_\delta = \inf \left\{ \sum_i \delta(p^{n_i}) : u = \sum_i e_i p^{n_i}, e_i \in \{1, -1\}, n_i \in \mathbf{N} \right\}.$$

We denote by $\mathbf{Z}_\delta$ the topological group $\mathbf{Z}$ with the metric induced by $\|\cdot\|_\delta$. This topology is strictly between the p -adic topology and the strongest topology.

For distinct primes p, q , we construct $\delta : \{p^n : n \in \mathbf{N}\} \rightarrow \mathbf{R}$ and $\epsilon : \{q^n : n \in \mathbf{N}\} \rightarrow \mathbf{R}$ with certain property. We need another corollary to Theorem 3.1.

Lemma 5.1. *We consider the equation $x_1 + \cdots + x_k = y_1 + \cdots + y_l$ in Lemma 3.2 under the restriction such that no (non-empty) subsum of $x_1 + \cdots + x_k$ or of $y_1 + \cdots + y_l$ vanishes, in particular $x_i \in U_S, y_j \in U_T$. Then the number of solutions is finite.*

Proof. We proceed similarly as in the proof of Lemma 3.2. We change the definition of I -solutions according to the restriction: they are tuples $\langle x_i : i \in I \rangle$ such that $\sum_{i \in I} x_i = 0$, $\{x_i : 1 \leq i \leq k, i \in I\} \subseteq U_S$, $\{x_i : k+1 \leq i \leq k+l, i \in I\} \subseteq U_T$ and no subsum of $\sum_{1 \leq i \leq k, i \in I} x_i$ or of $\sum_{k+1 \leq i \leq k+l, i \in I} x_i$ vanishes. So there is no I -solutions in the degenerate case.

By the induction on the size of I , we prove the finiteness of I -solutions. Assume that I is not degenerate. Then there are two possibilities for an I -solution: either it has no vanishing non-trivial sums or it is the union of a J -solution and an $I \setminus J$ -solution for a non-trivial subset J of I . Hence Theorem 3.1 and the induction hypothesis yields the conclusion. $\square$

Suppose that a non-zero integer u has representations as $u = \sum_{1 \leq i \leq k} e_i p^{m_i} = \sum_{1 \leq j \leq l} f_j q^{n_j}$ with $e_i, f_j \in \{1, -1\}$ and $m_i, n_j \in \mathbf{N}$. To estimate $\|u\|_\delta$ and $\|u\|_\epsilon$, we may assume that no subsum is zero. Due to the lemma above, for fixed k, l , there are only finitely many summands p^{m_i} which appear in such representations for some non-zero integer u . Let $F(p, q, k, l)$ denote the finite set of such p^{m_i} . For a positive integer s , set $F(p, q, s) = \bigcup_{k, l \leq s} F(p, q, k, l)$. Then it is a finite set non-decreasing with respect to s and $\bigcup_{s \in \mathbf{N}} F(p, q, s) = \{p^n : n \in \mathbf{N}\}$.

Now we define particular δ and ϵ by

$$\delta(p^n) = 1/\min\{s : p^n \leq \max F(p, q, s)\},$$

$$\epsilon(q^n) = 1/\min\{s : q^n \leq \max F(q, p, s)\}.$$

Then they are non-increasing and tend to 0. Moreover we have

$$p^n \in F(p, q, s) \Rightarrow \delta(p^n) \geq 1/s,$$

$$q^n \in F(q, p, s) \Rightarrow \epsilon(q^n) \geq 1/s.$$

We estimate $\|u\|_\delta$ and $\|u\|_\epsilon$ for a integer u with the representations above. If we set $s = \max\{k, l\}$, then $\delta(p^{m_i}), \epsilon(q^{n_j}) \geq 1/s$ for each summand. Without loss of generality, we may assume $k \geq l$. Then we have

$$\sum_{1 \leq i \leq k} \delta(p^{m_i}) \geq k \cdot (1/s) = 1,$$

and hence $\|u\|_\delta \geq 1$.

Accordingly it does not occur that both $\|u\|_\delta$ and $\|u\|_\epsilon$ are less than 1 for a non-zero integer u . Therefore the diagonal is discrete in $\mathbf{Z}_\delta \times \mathbf{Z}_\epsilon$.

Theorem 5.2. *Neither $X(1)$ nor $X(2)$ is not preserved under the product $\mathbf{Z}_\delta \times \mathbf{Z}_\epsilon$ for δ and ϵ decreasing slowly enough.* $\square$

REFERENCES

- [1] A.V. Arhangel'skiĭ, Frequency spectrum of a topological space and the classification of spaces (Russian), Dokl. Akad. Nauk. SSSR 206 (1972), 265–268.
- [2] A.V. Arhangel'skiĭ, The spectrum of frequencies of a topological space and the product operation (Russian), Trudy Moskov. Mat. Obshch. 40 (1979), 171–206.
- [3] R. Brown, P.J. Higgins and S.A. Morris, Countable products and sums of lines and circles: their closed subgroups, quotients and duality properties, Math. Proc. Cambridge Philos. Soc. 78 (1975), 19–32.
- [4] S. Kaplan, Extension of Pontryagin duality II: direct and inverse sequences, Duke Math. J. 17 (1950), 419–435.
- [5] N. Noble, k -groups and duality, Trans. Amer. Math. Soc. 151 (1970), 551–561.

- [6] P.J. Nyikos, Metrizable and the Fréchet-Urysohn property in topological groups, Proc. Amer. Math. Soc. 83 (1981), 793–801.
- [7] I.V. Protasov and E.G. Zelenyuk, Topologies on groups determined by sequences, Math. Studies Monograph Ser., Vol. 4, VNTL Publishers, L'viv, 1999.
- [8] W.M. Schmidt, Diophantine approximations and Diophantine equations, Lecture Notes Math., Vol. 1467, Springer-Verlag, 1991.
- [9] A. Shibakov, Examples of sequential topological groups under the continuum hypothesis, Fund. Math. 151 (1996), 107–120.
- [10] S. Todorčević, Some applications of S and L combinatorics, (F. D. Tall ed.) The work of Mary Ellen Rudin, Ann. New York Acad. Sci. 705 (1993), 130–167.
- [11] N.Th. Varopoulos, Studies in harmonic analysis, Proc. Cambridge Philos. Soc. Math. Phys. Sci. 60 (1964), 465–516.
- [12] E.G. Zelenyuk and I.V. Protasov, Topologies on abelian groups (Russian), Izv. Akad. Nauk SSSR Ser. Mat. 54 (1990), 1090–1107, translation in Math. USSR-Izv. 37 (1991), 445–460.

EDA LABORATORY, SCHOOL OF SCIENCE AND ENGINEERING, WASEDA UNIVERSITY, SHINJUKU-KU, TOKYO, 169-8555, JAPAN

E-mail address: `higasik@logic.info.waseda.ac.jp`